


HUMAN SPORT

MEDICINE 2020 Vol. 20 No. S1



ISSN 2500-0209 (Print)
ISSN 2500-0195 (Online)

South Ural State University

Main objectives of the journal are:

- to disseminate information about scientific research and development performed by scientists and top specialists with data distribution within the Russian Federation, CIS countries and far-abroad countries;
- to develop scientific fields and schools round the journal;
- to provide information support of priority scientific studies;
- to popularize advanced scientific ideas;
- to conduct propaganda of modern scientific physiological and medical technologies including recuperation;
- to conduct assessment of functional and metabolic condition, simulation and prediction in sports development on the basis of supercomputer technologies;
- to develop methodological provisions associated with foregoing areas and to implement them.

Editorial Board:

Isaev A.P. (*Chief Editor*), Honored Worker of Science, Dr. of Sci. (Biol.), Prof., South Ural State University, Chelyabinsk, Russian Federation;

Erlikh A.V. (*Deputy Chief Editor*), Dr. of Sci. (Biol.), Prof., South Ural State University, Chelyabinsk, Russian Federation;

Nenasheva A.V. (*Executive Secretary*), Dr. of Sci. (Biol.), Prof., South Ural State University, Chelyabinsk, Russian Federation;

Ushakov A.S. (*technical secretary*), South Ural State University, Chelyabinsk, Russian Federation

Editorial Council:

Schurova E.N., Dr. of Sci. (Biol.), Russian Ilizarov Scientific Center for Restorative Traumatology and Orthopaedics, Kurgan, Russian Federation;

Abzalov R.A., Dr. of Sci. (Biol.), Institute of Fundamental Medicine and Biology, Kazan (Volga Region) Federal University, Kazan, Russian Federation;

Pavlova V.I., Dr. of Sci. (Biol.), Chelyabinsk State Pedagogical University, Chelyabinsk, Russian Federation;

Sashenkov S.L., Dr. of Sci. (Med.), South Ural State Medical University, Chelyabinsk, Russian Federation;

Kuznetsov A.P., Honored Worker of Science, Dr. of Sci. (Biol.), Prof., Kurgan State University, Kurgan, Russian Federation;

Iryanov Y.M., Dr. of Sci. (Biol.), Prof., Russian Ilizarov Scientific Center for Restorative Traumatology and Orthopaedics, Kurgan, Russian Federation;

Vazhenin A.V., Academician of the Russian Academy of Sciences, Dr. of Sci. (Med.), South Ural State Medical University, Chelyabinsk, Russian Federation;

Yushkov B.G., Dr. of Sci. (Med.), Prof., Institute of Immunology and Physiology of the Russian Academy of Sciences, Ekaterinburg, Russian Federation;

Manukhina E.B., Dr. of Sci. (Med.), Health Science Center of the University of North Texas, USA;

Gubin A.V., Dr. of Sci. (Med.), Prof., Russian Ilizarov Scientific Center for Restorative Traumatology and Orthopaedics, Kurgan, Russian Federation;

Mileva K., PhD (Biomed. Eng.), Research Center of Sports Science and Nutrition, London South Bank University, Great Britain;

Shlyk N.I., Honored Science Worker of the Udmurt Republic, Dr. of Sci. (Biol.), Prof., Udmurt State University, Izhevsk, Russian Federation;

Escobar-Molina R., Prof., University of Granada, Spain;

Mazin H.K., PhD, University of Babylon, Iraq;

Nikityuk D.B., Dr. of Sci. (Med.), Professor, Russian Academy of Sciences, Moscow, Russian Federation;

Le Panse B., PhD, Le Panse Academy, Paris, France;

Poznyakovsky V.M., Dr. of Sci. (Biol.), Merited Scientist, Prof., Kemerovo Agrarian University, Kemerovo, Russian Federation;

Chereshnev V.A., Dr. of Sci. (Med.), member of the Russian Academy of Sciences, Professor, Ekaterinburg, Russian Federation;

Waluch K., PhD, associate professor of the department for teaching in Pawel Wlodkowica University College, Plock; member of TEAM EUROPE in the European Commission and Plock scientific society, Institute for Sport and Education Development, Warsaw, Poland;

Sonkin V.D., Dr. of Sci. (Biol.), Russian Academy of Education, Russian State University of Physical Education, Sport, Youth and Tourism, Moscow, Russian Federation;

Erlikh V.V., Dr. of Sci. (Biol.), Prof., South Ural State University, Chelyabinsk, Russian Federation;

Cherepov E.A., Doctor of Pedagogical Sciences, Associate Professor, South Ural State University, Chelyabinsk, Russian Federation;

Talaghir L.G., PhD, Prof., "Dunarea de Jos" University of Galati, Galati, Romania;

Dopsaj M., PhD, Prof., University of Belgrade, Belgrade, Serbia;

Bendíková E., PhD, Ass. Prof. Paed Dr., Matej Bel University, Banská Bystrica, Slovak Republic;

Kapilevich L.V., Dr. of Sci. (Med.), Prof., National Research Tomsk State University, Tomsk, Russia

CONTENTS

PHYSIOLOGY

RADZHABKADIEV R.M., VYBORNAYA K.V., LAVRINENKO S.V., VASILEV A.V. Thyroid Status of Athletes in Various Disciplines	5
STRELYAEVA O.P., SHAYAKHMETOVA E.Sh., SHURUKHINA G.A., MATVEEVA L.M., VALITOVA A.I. Features of Autonomic Regulation of Heart Rate in University Teachers When Using Preventive Measures	13
BOBKOVA S.N., NALOBINA A.N., ZVEREVA M.V., FEDOROVA E.Yu., ISKAKOVA Zh.T. Pharmaceutical Provision and Physical Activity for Supporting Working Capacity of Military and Security Services	20

SPORTS TRAINING

EVSEEV S.P., EVSEEVA O.E., AKSENOV A.V., Vishnyakova Yu.Yu., SHELEKHOV A.A. All-Russian Physical Education and Sports Program "Ready for Labor and Defense" (GTO) for Disabled People and Persons with Health Limitations: Concept, State, Prospects for Development	27
YAKHIN D.Kh., BELENKOV A.S., EPISHEVA A.A., SUMAK E.N., YAKHINA T.V. Issues of Masters Sport and the Ways to Solve Them (Based on the Example of Track-and-Field)	36
VAVILOV A.L., ANFILATOVA O.V., LAPTEV A.V., SIRAKOVSKAYA Ya.V. Identification of Physical Fitness Tests for Basketball Players	46
GÜMÜŞ H., GENÇOĞLU C. PlayerLoad tm and Heart Rate Response to Small-Sided Games Specialized to Additional Field Player Rule in Handball	55
SEDOCHENKO S.V., CHERNYKH A.V., Savinkova O.N. Stabilometric Parameters of Balance Development in Athletes Through Biological Feedback	62
SEVDALEV S.V., SKIDAN A.A., VRUBLEVSKIY E.P. Organizational and Methodical Aspects of Individualization of Health Improving Female Training	69
NIKOLAEV E.A., SURIKOVA N.V., CHUPROVA E.D., GOMBOEV B.B., UVAROV V.A. Justification of Hiking Distance as a Kind of Test Within the All-Russian Physical Education and Sports Training Program "Ready for Labour and Defence" for Girls	77
IZOV N., ALEXANDROVA A., PETROV L., KACHAUNOV M., SHEYTANOVA T., KOLIMECHKOV S. Dynamics of Training Distress, Performance, and Excretion of Cortisol and Cortisone in Urine During Six Weeks of Training in Elite Swimmers	84
BOGDANOV I.V., RYCHKOVA N.A. Post-Training and Recovery Activity in the Continuous Physical Training of Young People in the Kola Polar Region	92
POMERANTSEV A.A. The Features of Perceptual Processes During Qualitative Biomechanical Analysis ..	98
OSIPOV A.Yu., GURALEV V.M., NAGOVITSYN R.S., KAPUSTIN A.G., KOVYAZINA G.V. Justification of Using Crossfit Training in Judo	109

SPORT NUTRITION

NAUMENKO N.V., Potoroko I.Yu., KALININA I.V., NENASHEVA A.V., BOTVINNIKOVA V.V. Possibilities of Regulating Stress-Protective Activity of Food Products to Improve Immunity under the Conditions of the COVID-19 Pandemic	116
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----

REHABILITATION AND SPORT MEDICINE

LITOVCHENKO O.G., MAXIMOVA A.S. Psychophysiological Features of Volleyball Players with Hearing Impairment	128
KORKMAZOV M.Yu., KORKMAZOV A.M., DUBINETS I.D., SMIRNOV A.A., KORNOVA N.V. Influence of Non-Drug Therapy on Rehabilitation Time and Skeet Shooting after Rhinosurgical Interventions	136

DYNAMICS OF TRAINING DISTRESS, PERFORMANCE, AND EXCRETION OF CORTISOL AND CORTISONE IN URINE DURING SIX WEEKS OF TRAINING IN ELITE SWIMMERS

N. Iзов¹, rector@nsa.bg, ORCID: 0000-0002-1362-5291,
 A. Alexandrova¹, a_alexandrova_bas@yahoo.com, ORCID: 0000-0002-7007-3665,
 L. Petrov¹, dr.lubomir.petrov@gmail.com, ORCID: 0000-0003-1209-959X,
 M. Kachaunov¹, mihail.kachaunov@gmail.com, ORCID: 0000-0002-6877-6054,
 T. Sheytanova¹, tanyasheytanova@mail.bg, ORCID: 0000-0003-0906-6834,
 S. Kolimechkov^{1, 2}, dr.stefan.kolimechkov@gmail.com, ORCID: 0000-0003-0112-2387

¹National Sports Academy, Sofia, Bulgaria,

²STK Sport, London, Great Britain

Aim. Optimal balance between training intensity and recovery is of particular importance for elite swimmers in order to improve their results. The aim of this study was to record the dynamics of training distress, performance, and excretion of cortisol and cortisone in urine during six weeks of training in elite swimmers. **Materials and Methods.** Twenty-four participants (10 women and 14 men) from the national swimming team of Bulgaria took part in this study, with an average age of 18.7 ± 3.78 years. Training distress (TDS) and urine concentration of cortisol and cortisone were evaluated every two weeks. In total, the measurements were taken on the 1-st (T1), 14-th (T2), 28-th (T3) and 42-nd days (T4). Anthropometric measurements were also taken at T1 and T4, and body fat percentage and muscle mass percentage were calculated by skinfold methods. **Results.** The TDS score at T4 (6.92 ± 7.15) was significantly lower than this at T1 (14.96 ± 10.63) and T2 (15.21 ± 12.44). The concentrations of cortisol at T3 (35.9 ± 47.7) and T4 (35.0 ± 24.2) were also significantly lower than that registered at T1 (82.7 ± 62.8). The concentration of cortisone did not show any significant differences across T1 to T4, but the sum of cortisol and cortisone urine concentration was significantly lower in T3 and T4 vs T1. **Conclusion.** The significantly reduced cortisol concentration in urine in T4, as well as the sum of the concentrations of cortisol and cortisone, were in line with the reduction of the TDS score in T4.

Keywords: cortisol, cortisone, swimmers, overtraining, training distress score.

Introduction

Elite swimmers perform large amounts of training at high intensity to improve their competition results. In order to avoid the development of overtraining syndrome, they have to maintain an optimal balance between training intensity and recovery [1].

Studies on urinary cortisol/cortisone ratio (C/Cn ratio) overnight [5] and C/Cn ratio for 24 h [1] have been conducted to monitor the effect of the training load on elite swimmers. The hypothalamic-pituitary-adrenal axis (HPA axis) plays the main role in the adaptation for endurance training and for the response to intensive exercises. Repeated exercises bring adaptation of the HPA axis with decreased sensitivity of tissues to glucocorticoids [6] and reduced glucocorticoids receptor levels [3]. Urine sampling methods, as opposed to plasma, are applied to assess HPA axis responses when athletes perform large amounts of training and participation in competitions, without any disturbances to their training

conditions. These methods are non-invasive, non-stressful, reliable, and practical [1]. Plasma cortisol levels are modulated by variations of their binding protein (CBG) and bear a low level of correlation with the rate of cortisol production, unless differences in CBG are corrected, whereas urinary free cortisol concentrations are independent of CBG concentrations and thus, closely reflect the production of 24-h free active plasma cortisol [2].

Different physiological variables have been investigated as a potential marker of training distress, which include metabolic, cardiovascular, immunological, neuromuscular, and endocrinological measures [8]. However, due to their high intra-individual variability, those variables have shown only weak relationships with training distress [7, 8]. On the other hand, psychological and behavioural measures have been observed to have stronger relationships. Additionally, they are efficient, inexpensive and non-invasive [16].

Grove et al. published findings from three

experimental studies which estimated the short-term training distress and performance readiness in swimmers [7]. The authors confirmed that reductions in performance capabilities which are induced by the training workload were associated with parallel increase in the score of the Training Distress Scale (TDS). TDS is a 19-item questionnaire which measures generated distress symptoms previously identified by Fry et al. [4]. The link between TDS scores and swimming performance was evident from the results of the swimmers with low levels of distress symptoms 2 weeks before a major competition, who swam better during the event than those who reported higher levels of distress symptoms [7].

Aim

The aim of this study was to record the dynamics of training distress, performance, and excretion of cortisol and cortisone in urine during six weeks of training in elite swimmers.

Materials and Methods

Participants

Twenty-four elite swimmers (10 women and 14 men) from the national swimming team of Bulgaria took part in this study. They had an average age of 18.7 ± 3.78 years.

Design of the study

The athletes were tested over a 6-week period starting 2 weeks after the National Team Championship and finishing 2 weeks before the National Individual Championship (both in a 50 m swimming pool). The degrees of training distress and urine concentration of cortisol and cortisone were evaluated every two weeks. In total, the measurements were taken four times: on the 1-st (T1), 14-th (T2), 28-th (T3) and 42-nd days (T4) on Monday mornings. Additionally, anthropometric measurements were conducted at T1 and T4.

Training programme

The study was conducted in the summer sea-

son (3rd macrocycle) stretching from April to July, in which the training sessions in a 50 m swimming pool were performed. The reported six-week period was part of the general preparation phase, and it started right after the 2 weeks of recovery phase following the main competition. The swimmers trained twice per day from Monday to Friday, performing a total of 10 training sessions a week. During this period, the training process of the swimmers is associated with a gradually increased training volume, starting from 40 km for the first week and finishing at 65 km for the last. The main goals for this training phase are to improve aerobic capacity and to maintain anaerobic power and speed. The training has to produce central circulatory and respiratory adaptations which improve the delivery of oxygen and nutrients in the muscles [14], and as a result of that, an increased $VO_2\max$ is expected. The training programme of the participants in this study, including training volume (km) and training zones (% of all volume), is presented in Table 1.

HR-bpm

Anthropometry

The following anthropometric measurements were taken: height, weight and arm span. Body fat percentage (Fat%) and muscle mass percentage (MM%) were calculated by using skinfold methods [13–15].

Training distress

The Training Distress Scale (TDS) [7] was translated into Bulgarian and applied to assess the athletes' degree of training distress and performance readiness. The TDS consists of 19 items which measure the distress symptoms previously identified by Fry et al. [4]. The responses were given by using a 5-point bipolar scale extending from 'not at all (0)' to 'extreme amount (4)'. The final score was calculated as the sum of all points.

Table 1

The six-week training programme of the elite swimmers

Week	1	2	3	4	5	6
Training zones/Volume (km)	40	45	50	55	60	65
I. Basic Endurance HR 120-150	59 %	59 %	59 %	50 %	50 %	50 %
II. Threshold Endurance HR 150-180	25 %	25 %	25 %	25 %	25 %	25 %
III. High performance endurance HR >180	4 %	4 %	4 %	8 %	8 %	8 %
IV. Anaerobic: Lactate tolerance + Lactate production + Race pace training	2 %	2 %	2 %	4 %	4 %	4 %
V. Sprint	3 %	3 %	3 %	3 %	3 %	3 %
VI. Recovery HR < 120	7 %	7 %	7 %	10 %	10 %	10 %

Спортивная тренировка

Additionally, first-morning urine samples were taken to determine the concentration of cortisol and cortisone. The participants, with only 3 exceptions, noted that this was their only urine for the whole night. The competitors always had 60 hours of recovery (from Friday night to Monday morning) before the samples were taken.

Chemicals and reagents

Formic acid (HPLC grade) was obtained from Merck (Darmstadt, Germany). Acetonitrile, methanol and tert-butyl methyl ether (HPLC grade), K₂CO₃ and KHCO₃ puriss p.a. were purchased from Sigma-Aldrich (Steinheim, Germany). Cortisol and cortisone were from LGC (Luckenwalde, Germany). D₃-Methyltestosterone was from RIKILT (Wageningen, Netherlands).

Sample preparation

Two millilitres by volume of urine were mixed with 0.5 mL carbonate buffer pH 9.6 (K₂CO₃/KHCO₃ 20/20, w/w) and shaken. After the addition of 20 µL internal standard (ISTD, d₃-methyltestosterone, 200 ng/mL), the urine sample was extracted by 3.0 mL of t-butyl-methyl ether for 20 min and centrifuged at 5000 rpm for 10 min. The aqueous phase was frozen and the organic layer was isolated in another tube and evaporated to dryness at 55 °C under a stream of nitrogen. The residue was reconstituted in 200 µL of CH₃OH/ 0.2% formic acid, H₂O (10:90).

A five-point calibration curve and two quality controls (QC) in water (Table 2) were used for the quantification of samples.

Instrumentation

LC-MS-MS analysis was performed on an Agilent Technologies UHPLC 1290 Infinity with 6460 Triple Quad mass spectrometer.

A reversed-phase ZORBAX Eclipse Plus C18 Rapid Resolution HD column (2.1 × 50 mm 1.8 Micron) was used for the analyses. The mobile phase was composed of 0.2% formic acid, H₂O as solvent A and acetonitrile as solvent B. A gradient (Table 3) was run at 0.4 mL/min and the injected volume was 10 µL.

MS/MS detection was performed in multiple reaction monitoring (MRM) mode and electrospray ionization (ESI) in the positive ion mode. Acquisition parameters were summarised in Table 4.

The obtained sample concentrations were adjusted to a urine specific gravity (SG) of 1.020 based on the following equation:

$$\text{Conc}_{\text{corrected}} = \text{Conc}_{\text{measured}} \times (1.020 - 1) / (\text{SG} - 1).$$

Statistical analyses

Statistical analysis was performed using the GraphPad Prism 5.0 software. The significance of differences of anthropometric mean values was evaluated by paired Student's t-test. Kolmo-

Table 2

LOD, LOQ, calibration level and QC for method performance in ng/mL

Substance	LOD, ng/mL	LOQ, ng/mL	cal-1, ng/mL	cal-2, ng/mL	cal-3, ng/mL	cal-4, ng/mL	cal-5, ng/mL	QC-1, ng/mL	QC-2, mg/mL
Cortisol	0,186	0,400	0,400	10,000	40,000	100,000	200,000	22,000	62,000
Cortisone	0,135	0,400	0,400	10,000	40,000	100,000	200,000	22,000	60,000

Table 3

Liquid chromatography solvent gradients for the UHPLC MS/MS method

Time (min)	0	1	1.5	3	4	5.5	6	8	8.7	9	9.5	9.7	12	12.2	17
A (%)	95	95	90	80	70	70	60	60	30	20	20	5	5	95	95
B (%)	5	5	10	20	30	30	40	40	70	80	80	95	95	5	5

Table 4

Multiple Reaction Monitoring (MRM) and mass spectrometer conditions (ESI)

Substance	RT, min	Precursor Ion	Product Ion	CE, eV	Polarity
Cortisol	4,88	363,22	121,1	23	Positive
		363,22	105,1	51	Positive
Cortisone	4,92	361,20	163,1	23	Positive
		361,20	121,1	35	Positive
		361,20	105,1	47	Positive
d ₃ -MT	7,60	306,20	109,1	20	Positive
		306,20	97,1	20	Positive

gorov-Smirnov test of normality was performed for TDS, Cortisol (C), Cortisone (Cn) and C/Cn ratio. The differences between the obtained values for each measurement were evaluated by the non-parametric Friedman's test for repeated measures with Dunns *post hoc* test for the significance of differences between each pair. The data is presented as Mean \pm Standard Error (SE) in graphics and as Mean \pm Standard Deviation (SD) in the text.

Results

The anthropometric parameters which were measured in T1 and T4 are presented in Fig. 1.

There were no statistically significant differences in body composition, both in males and in females.

The analysis of the responses from the Training Distress Scale (TDS) showed a good internal consistency for the composite measure, with an average of Cronbach's alpha of 0.871 across administrations. The TDS score dynamics are presented in Fig. 2. The TDS score at T4 (6.92 ± 7.15) was significantly lower than that at T1 (14.96 ± 10.63), as well as that at T2 (15.21 ± 12.44). T1 14.96 ± 10.63 ; T2 15.21 ± 12.44 ; T3 11.54 ± 7.91 ; T4 6.92 ± 7.15 .




Fig. 1. Anthropometric parameters of female (Left) and male (Right) swimmers at T1 and T4




Fig. 2. Dynamics of TDS score for the elite swimmers across T1 to T4:

* p < 0.05 vs T4; *** – p < 0.001 vs T4




Fig. 3. Dynamics of cortisol and cortisone urine concentration across T1 to T4:
* $p < 0.05$ vs T1




Fig. 4. Dynamics of the cortisol/cortisone ratio across T1 to T4




Fig. 5. Dynamics of the sum of cortisol and cortisone urine concentration across T1 to T4:
* $p < 0.05$ vs T1

The dynamics of cortisol and cortisone urine concentration are presented in Fig. 3. The concentration of cortisol at T1 (82.7 ± 62.8) was significantly higher than those at T3 (35.9 ± 47.7) and T4 (35.0 ± 24.2). The concentration of cortisone did not show any significant differences at any of the times tested (T1 to T4).

The dynamics of cortisol/cortisone ratio (C/Cn ratio) are presented in Fig. 4, and that of the sum of cortisol and cortisone urine concentration is presented in Fig. 5. The C/Cn ratio did not show any significant differences at any of the times tested (T1 to T4), while the sum of cortisol and cortisone urine concentration was significantly lower in T3 and T4 vs T1.

Discussion

The body composition data of the elite swimmers during the investigated period remained the same, which suggested that the nutritional status of the participants was unchanged. As a side observation, a bigger difference between the height and the arm span, more visible in men (about 7 cm), was registered in this study. Similar findings were reported for swimmers in the literature [12].

The adapted Training Distress Scale (TDS) has been shown to be an easy and appropriate instrument which can be used in everyday coaching practice, and as a tool for longitudinal studies. The results from our study showed that the TDS scores decreased during the investigated period (Fig. 2), despite the training volume increasing by 5 km per week (Table 1). This was also backed up by the reduced concentration of cortisol in T3 and T4 vs T1.

In our study, in almost all cases the morning urine was the all-night urine, and the concentration of the hormones measured was adjusted to its density. Thus, the effect of a volume of urine was considerably reduced, and the measured cortisol and cortisone concentrations reflected the total amount of hormones released overnight.

Conclusions

The significantly reduced cortisol concentration in urine in T4, as well as the sum of the concentrations of the cortisol and cortisone, are in line with the reduction of the TDS score in T4. The lower concentration of the stress hormone, cortisol, in the urine, and the lack of subjective symptoms of training distress led to good physical and mental recovery of the athletes after 60 hours of rest.

Acknowledgements. This study was part of the project “Non-invasive monitoring of recovery and effects of glutathione supplementation on

elite swimmers” supported by Grant ЦПАК 12/15.02.2018 by the National Sports Academy, Sofia, Bulgaria.

References

1. Atlaoui D., Duclos M., Gouarne C. et al. The 24-h Urinary Cortisol/Cortisone Ratio for Monitoring Training in Elite Swimmers. *Medicine and Science in Sports and Exercise*, 2004, vol. 36 (2), pp. 218–224. DOI: 10.1249/01.MSS.0000113481.03944.06
2. Bright G.M. Corticosteroid-Binding Globulin Influences Kinetic Parameters of Plasma Cortisol Transport and Clearance. *J Clin Endocrinol Metab*, 1995, vol. 80 (3), pp. 770–775. DOI: 10.1210/jc.80.3.770
3. Duclos M., Gouarne C., Bonnemaïson D. Acute and Chronic Effects of Exercise on Tissue Sensitivity to Glucocorticoids. *J Appl Physiol*, 1985, vol. 94 (3), pp. 869–875. DOI: 10.1152/jappphysiol.00108.2002
4. Fry R.W., Grove J.R., Morton A.R. et al. Psychological and Immunological Correlates of Acute Overtraining. *Br J Sports Med*, 1994, vol. 28 (4), pp. 241–246. DOI: 10.1136/bjism.28.4.241
5. Gouarne C., Groussard C., Gratas-Delamarche A. et al. Overnight Urinary Cortisol and Cortisone Add New Insights Into Adaptation to Training. *Medicine and Science in Sports and Exercise*, 2005, vol. 37 (7), pp. 1157–1167. DOI: 10.1249/01.mss.0000170099.10038.3b
6. Grasso G., Lodi L., Lupo C., Muscettola M. Glucocorticoid Receptors in Human Peripheral Blood Mononuclear Cells in Relation to Age and to Sport Activity. *Life Sci*, 1997, vol. 61 (3), pp. 301–308. DOI: 10.1016/S0024-3205(97)00386-X
7. Grove J.R., Main L.C., Partridge K. et al. Training Distress and Performance Readiness: Laboratory and Field Validation of a Brief Self-Report Measure. *Scand J Med Sci Sports*, 2014, vol. 24 (6), pp. 483–490. DOI: 10.1111/sms.12214
8. Halson S.L., Jeukendrup A.E. Does Overtraining Exist? An Analysis of Overreaching and Overtraining Research. *Sports Med*, 2004, vol. 34 (14), pp. 967–981. DOI: 10.2165/00007256-200434140-00003
9. Hug M., Mullis P.E., Vogt M. et al. Training Modalities: Over-Reaching and Over-Training in Athletes, Including a Study of the Role of Hormones. *Best Pract Res Clin Endocrinol Metab*, 2003, vol. 17 (2), pp. 191–209. DOI: 10.1016/S1521-690X(02)00104-5
10. Jackson A.S., Pollock M.L. Generalized

Equations for Predicting Body Density of Men. *Br J Nutr*, 1978, vol. 40 (3), pp. 497–504. DOI: 10.1079/BJN19780152

11. Jackson A.S., Pollock M.L., Ward A. Generalized Equations for Predicting Body Density of Women. *Medicine and Science in Sports and Exercise*, 1980, vol. 12 (3), pp. 175–181. DOI: 10.1249/00005768-198023000-00009

12. Latt E., Jurimae J., Maestu J. et al. Physiological, Biomechanical and Anthropometrical Predictors of Sprint Swimming Performance in Adolescent Swimmers. *J Sports Sci Med*, 2010, vol. 9 (3), pp. 398–404.

13. Mackinnon L.T., Hooper S.L., Jones S. et al. Hormonal, Immunological, and Hemato-

logical Responses to Intensified Training in Elite Swimmers. *Medicine and Science in Sports and Exercise*, 1997, vol. 29 (12), pp. 1637–1645. DOI: 10.1097/00005768-199712000-00014

14. Maglischo E. *Swimming Fastest: Human Kinetics*. 2003.

15. Martin A.D., Spent L.F., Drinkwater D.T., Clarys J.P. Anthropometric Estimation of Muscle Mass in Men. *Medicine and Science in Sports and Exercise*, 1990, vol. 22 (5), pp. 729–733. DOI: 10.1249/00005768-199010000-00027

16. O'Connor P. Overtraining and Staleness. In W. Morgan (Ed.), *Physical Activity and Mental Health*. Washington: Taylor & Francis. 1997, pp. 145–160.

Received 15 March 2020

УДК 797.2.015

DOI: 10.14529/hsm20s111

ДИНАМИКА ТРЕНИРОВОЧНОГО ДИСТРЕССА, ПРОИЗВОДИТЕЛЬНОСТИ И УРОВНЯ КОРТИЗОЛА И КОРТИЗОНА В МОЧЕ У ЭЛИТНЫХ ПЛОВЦОВ ВО ВРЕМЯ ШЕСТИНЕДЕЛЬНОГО ТРЕНИРОВОЧНОГО ЦИКЛА

Н. Изов¹, А. Александрова¹, Л. Петров¹, М. Качаунов¹, Т. Шейтанова¹, С. Колимечков^{1, 2}

¹Национальная Спортивная Академия, г. София, Болгария,

²Компания STK Sport, г. Лондон, Великобритания

Цель. Оптимальный баланс между интенсивностью тренировок и восстановлением особенно важен для улучшения результатов элитных пловцов. Данное исследование было нацелено на выявление динамики тренировочного дистресса, производительности и экскреции кортизола и кортизона у элитных пловцов во время шестинедельного тренировочного цикла. **Материалы и методы.** В исследовании приняли участие 24 спортсмена (10 женщин и 14 мужчин) национальной сборной Болгарии по плаванию, средний возраст которых составил $18,7 \pm 3,78$ года. Тренировочный дистресс и концентрация кортизола и кортизона в моче оценивались каждые две недели. Измерения проводились в 1-й (Т1), 14-й (Т2), 28-й (Т3) и 42-й дни (Т4). Антропометрические измерения проводились в 1-й и 42-й дни. Процентное содержание жира и мышечной массы оценивалось с использованием метода измерения толщины кожной складки. **Результаты.** Показатель тренировочного дистресса на 42-й день ($6,92 \pm 7,15$) был значительно ниже, чем в 1-й ($14,96 \pm 10,63$) и 14-й дни измерений ($15,21 \pm 12,44$). Концентрация кортизола на 28-й ($35,9 \pm 47,7$) и 42-й день ($35,0 \pm 24,2$) также была значительно ниже, чем в 1-й день измерений ($82,7 \pm 62,8$). Концентрация кортизона не показала каких-либо существенных различий между 1-м и 42-м днем измерений, однако сумма концентраций кортизола и кортизона в моче была значительно ниже на 28-й и 42-й дни по сравнению с 1-м днем измерений. **Заключение.** Значительно сниженная концентрация кортизола в моче на 42-й день измерений, а также сумма концентраций кортизола и кортизона соответствовали уменьшению показателя тренировочного дистресса к 42-му дню тренировочного цикла.

Ключевые слова: кортизол, кортизон, пловцы, перетренированность, показатель тренировочного дистресса.

Николай Изов, PhD, ректор, профессор, кафедра плавания, Национальная спортивная академия. Болгария, 1700, г. София, р-он Студентский град. E-mail: rector@nsa.bg, ORCID: 0000-0002-1362-5291.

Альбена Александрова, PhD, доцент, кафедра физиологии и биохимии, Национальная спортивная академия. Болгария, 1700, г. София, р-он Студентский град. E-mail: a_alexandrova_bas@yahoo.com, ORCID: 0000-0002-7007-3665.

Любомир Петров, PhD, доцент, кафедра физиологии и биохимии, Национальная спортивная академия. Болгария, 1700, г. София, р-он Студентский град. E-mail: dr.lubomir.petrov@gmail.com, ORCID: 0000-0003-1209-959X.

Михаил Качаунов, PhD, соискатель, старший преподаватель, кафедра плавания, Национальная спортивная академия. Болгария, 1700, г. София, р-он Студентский град. E-mail: mihail.kachaunov@gmail.com, ORCID: 0000-0002-6877-6054.

Таня Шейтанова, PhD, соискатель, Национальная спортивная академия. Болгария, 1700, г. София, р-он Студентский град. E-mail: tanyasheytanova@mail.bg, ORCID: 0000-0003-0906-6834.

Стефан Колимечков, PhD, независимый зарегистрированный исследователь в Великобритании, компания STK Sport, г. Лондон, Великобритания, ул. Холсвилл-Роад 1, E16 1BE. E-mail: dr.stefan.kolimechkov@gmail.com, ORCID: 0000-0003-0112-2387.

Поступила в редакцию 15 марта 2020 г.

ОБРАЗЕЦ ЦИТИРОВАНИЯ

Dynamics of Training Distress, Performance, and Excretion of Cortisol and Cortisone in Urine During Six Weeks of Training in Elite Swimmers / N. Izov, A. Alexandrova, L. Petrov et al. // Человек. Спорт. Медицина. – 2020. – Т. 20, № S1. – С. 84–91. DOI: 10.14529/hsm20s111

FOR CITATION

Izov N., Alexandrova A., Petrov L., Kachaunov M., Sheytanova T., Kolimechkov S. Dynamics of Training Distress, Performance, and Excretion of Cortisol and Cortisone in Urine During Six Weeks of Training in Elite Swimmers. *Human. Sport. Medicine*, 2020, vol. 20, no. S1, pp. 84–91. DOI: 10.14529/hsm20s111